

The Angel Problem, Positional Games, and Digraph Roots

Strategies and Complexity

Dissertation zur Erlangung des Doktorgrades

vorgelegt am

Fachbereich Mathematik und Informatik
der Freien Universität Berlin

2004

von

Martin Kutz

Institut für Mathematik
Freie Universität Berlin
Arnimallee 2
14195 Berlin
Germany
kutz@math.fu-berlin.de

Betreuer und erster Gutachter:

Prof. Dr. Martin Aigner
Freie Universität Berlin
Arnimallee 2
D-14195 Berlin
Germany
`aigner@math.fu-berlin.de`

zweiter Gutachter:

Prof. Dr. Gyula O. H. Katona
Alfréd Rényi Institute of Mathematics,
Hungarian Academy of Sciences
H-1053 Budapest, Reáltanoda u. 13–15
Hungary
`ohkatona@renyi.hu`

Tag der Disputation: 7. Juli 2004

Preface

This thesis is about combinatorial games—mostly. It is also about graphs, directed graphs and hypergraphs, to a large extent; and it deals with the complexity of certain computational problems from these two areas. We study three different problems that share several of the above aspects, yet, they form three individual subjects and so we treat them independently in three self-contained chapters:

The angel-devil game. In the first chapter, we present improved strategies for an infinite game played on an infinite chess board, which has been introduced by Berlekamp, Conway, and Guy [8]. The *angel*, a chess person who jumps from square to square, tries to escape his opponent, the *devil*, who intends to strand the angel by placing obstacles on the board.

The open question about this game is, whether some angel who is allowed to make sufficiently large but bounded steps in each move, will be able to escape forever. Conway [11] has shown that certain quite natural escape attempts are bound to fail.

We attack this problem from the devil’s perspective, trying to improve upon a result from [8], which established that the ordinary chess king, who can be considered as an angel of minimal power, cannot escape. A reformulation of the game which focuses on the angel’s speed as the crucial quantity, allows us to show that certain faster “chess kings” can still be trapped. A second part of this chapter deals with angels on a three-dimensional board. We show that the new dimension grants the angel enough freedom to escape forever.

Weak positional games on hypergraphs. The games in the second chapter are very general versions of the well-known game of Tic-Tac-Toe. Two players alternately claim vertices of a hypergraph, the first player trying to get all vertices within some edge, his opponent striving to prevent this from happening.

Such *weak positional games* are known to be PSPACE-complete, but the respective hardness result from [39] utilizes edges of size up to 11. We analyze the restricted class of hypergraphs whose edges contain no more than three vertices each, trying to find optimal strategies for both players. We almost succeed. Under the additional restriction of almost-disjointness, that is, any two edges may share at most one vertex, we obtain a classification of such hypergraphs into those that yield a first player win and those who don’t, which immediately leads to efficiently computable optimal strategies for either player. Eventually, a new framework is introduced for describing values that individual parts of a hypergraph contribute to a game that is played on the whole hypergraph.

The complexity of digraph root computation. The final chapter is not about games. A k th root of a square Boolean matrix A is some other matrix R with $R^k = A$. Interpreting A as the adjacency matrix of a directed graph (digraph), we get an induced notion of powers for digraphs: the digraph D^k has an arc from a to b iff there is a walk of length exactly k from a to b in the digraph D .

The computational complexity of deciding whether a given Boolean matrix or, equivalently, a given digraph has a k th root, has been an open problem for twenty years. We answer this question by proving the problem NP-hard for every single integer $k \geq 2$. Our NP-completeness proof takes the graph-theoretic view, using basic concepts like paths, cycles, and vertex neighborhoods.

Besides the phenomena that make root finding hard, we discover a relation between digraph roots and graph isomorphism which materializes in form of an isomorphism-completeness result: For a special class of digraphs defined through arc subdivisions, root finding is of the same complexity as deciding whether two digraphs are isomorphic. This may come as a surprise since all problems known to be of this complexity are more or less obviously isomorphism problems. In abridged form, the results from this chapter have already appeared in [28].

Acknowledgments

I want to thank my advisor Martin Aigner. I am grateful for his support and guidance but also for granting me the freedom to find and pursue my own way. Many thanks to everybody at the mathematics department of Freie Universität Berlin who made the past three years such an amazing time. Especially to Mark de Longueville, Carsten Schultz, Stefan Geschke, Elmar Vogt, and Dennis Epple for endless talks about math and a thousand other things; and to Andrea Hoffkamp, simply for being there.

I thank the people from the computer-science department of Freie Universität Berlin, in particular Laura Heinrich-Litan for being supportive and especially Günter Rote for sharing his knowledge and for always being interested. Throughout my time at Freie Universität Berlin I was a member of the European graduate school “Combinatorics, Geometry, and Computation” supported by the Deutsche Forschungsgemeinschaft and I want to thank the faculty of the program for providing an excellent work environment. Also thanks to Manuel Bodirsky for fruitful cooperation, and to Vincenzo Marra and Jörg Zimmermann for long discussions.

I am grateful to the hospitable people at KAM in Prague, foremost to Martin Loeb, for his advice and his interest in my work; and to Emo Welzl’s group at ETH Zürich, in particular to Bernd Gärtner and Kaspar Fischer for a very productive time and to Ingo Schurr for sharing his office.

I also want to thank my teacher Arnold Schönhage, who influenced my mathematical thinking more than anybody else. Eventually, my thanks go to Sven Ehlert, Tim Bärmann, and Hans-Peter Jacobs for being very supportive; and to my girlfriend Petra Oyen for that and much more.

Berlin,
May 2004

Martin Kutz

Contents

Preface	iii
Acknowledgments	iv
Chapter 1. The Angel Problem	1
1. Angels, Kings, and Fools	1
2. From Finite to Infinite Games	5
3. The Need for Speed	6
4. Catching a $(2 - \epsilon)$ -King	12
5. An Escape into Space	19
Chapter 2. Weak Positional Games	29
1. Tic-Tac-Toe	29
2. Winning Ways	33
3. Decomposing Hypergraphs	37
4. Between the Docks	42
5. Playing for Breaker	55
6. Almost-Disjointness	62
7. Comparing Games	64
Chapter 3. Digraph Roots	71
1. Matrices and Digraphs, Powers and Roots	71
2. NP-Completeness	74
3. Roots and Isomorphism	77
Bibliography	89
Zusammenfassung	91

